

DATA NOTE

Open Access

# Non-targeted metabolomics and lipidomics LC–MS data from maternal plasma of 180 healthy pregnant women

Hemi Luan<sup>1,2</sup>, Nan Meng<sup>2</sup>, Ping Liu<sup>2</sup>, Jin Fu<sup>2</sup>, Xiaomin Chen<sup>2</sup>, Weiqiao Rao<sup>2</sup>, Hui Jiang<sup>2</sup>, Xun Xu<sup>2</sup>, Zongwei Cai<sup>1\*</sup> and Jun Wang<sup>2,3\*</sup>

## Abstract

**Background:** Metabolomics has the potential to be a powerful and sensitive approach for investigating the low molecular weight metabolite profiles present in maternal fluids and their role in pregnancy.

**Findings:** In this Data Note, LC–MS metabolome, lipidome and carnitine profiling data were collected from 180 healthy pregnant women, representing six time points spanning all three trimesters, and providing sufficient coverage to model the progression of normal pregnancy.

**Conclusions:** As a relatively large scale, real-world dataset with robust numbers of quality control samples, the data are expected to prove useful for algorithm optimization and development, with the potential to augment studies into abnormal pregnancy. All data and ISA-TAB format enriched metadata are available for download in the *MetaboLights* and *GigaScience* databases.

**Keywords:** Metabolomics, Lipidomics, Pregnancy, Metabolic phenotype, Maternal plasma

## Data description

### Purpose of collection

Metabolic variations occur during normal pregnancy to provide the developing foetus with a supply of nutrients required for development, and to ensure the health of the mother during gestation. The following dataset was collected to study metabolic phenotype variations in the maternal plasma, which are induced by pregnancy during each of its three trimesters [1]. All work described here is associated with this report, which used liquid chromatography–mass spectrometry (LC–MS) to survey the effects of pregnancy on the metabolite profiles of maternal plasma during pregnancy. A basic overview of the three datasets for the different experiments is shown in Figure 1. Full details of the batch numbers, run order and distribution of QCs can be seen in Additional file 1.

### Sample collection


A total of 180 healthy pregnant women were recruited from the Maternity and Child Health Hospital of Shenzhen, China. Full ethical approval for this study was provided by the Institutional Review Board of BGI-Shenzhen (No. 13045). All volunteers gave written informed consent and filled out an individual questionnaire at the time of collection for the study. Clinical information was obtained from neonatal and obstetrical medical records. Pregnancy was followed up to term to ensure participating women had normal term pregnancies and healthy babies. Individual participants were divided into six subgroups of 30 individuals according to their gestational weeks: class A (9w ~ 12w), class B (13w ~ 16w), class C (17w ~ 20w), class D (21w ~ 24w), class E (25w ~ 28w), and class F (29w ~ 40w). Weight, height, age and gestational week were recorded. Venous blood was collected in the morning before breakfast using VACU-ETTE® EDTA Blood Collection Tubes (Greiner Bio-One GmbH, NC, USA). Plasma samples were separated at 2200 g for 5 min at 4°C from vein blood and stored at

\* Correspondence: zwcail@hkbu.edu.hk; wangj@genomics.cn

<sup>1</sup>Department of Chemistry, Hong Kong Baptist University, 224 Waterloo Road, Kowloon Tong, Hong Kong, China

<sup>2</sup>BGI-Shenzhen, Building No.11, Beishan Industrial Zone, Yantian District, Shenzhen 518083, China

Full list of author information is available at the end of the article



–80°C until use. One sample was collected per patient, rather than multiple collections for each subject.

#### Non-targeted metabolic profiling spectral acquisition

Low molecular weight metabolites (<1000 Da) in the plasma samples were isolated using methanol precipitation with slight modifications to previously reported protocols [1,2]. To outline these briefly, 50 µL of thawed plasma was collected and precipitated using 200 µL of methanol. After centrifugation at 14000 *g* for 10 min at 4°C, the supernatant was transferred to 1.5 mL polypropylene tubes, and 10 µL of the supernatant injected into the LC–MS equipment. Quality control (QC) samples were also prepared by mixing equal volumes (10 µL) from each plasma sample before sample preparation as they were being aliquoted for analysis. This pooled sample was then used to estimate a mean profile representing all the analytes encountered during the analysis [3–4].

LC–MS data was acquired using a Shimadzu Prominence HPLC system (Shimadzu, Japan), coupled online to a LTQ Orbitrap Velos instrument (Thermo Fisher Scientific, MA, USA) set at a resolution of 30,000 (at *m/z* 400). Sample analysis was carried out in both positive and negative ion modes with the mass scanning range at 50–1500 *m/z* and capillary temperature of 350°C. Nitrogen sheath gas was set at a flow rate of 30 L/min, and auxiliary gas was set at a flow rate of 10 L/min. Spray voltage was 4.5 kV, with the LC–MS system run in binary gradient mode. Solvent A was 0.1% (v/v) formic acid/water and solvent B was 0.1% (v/v) formic acid/methanol; the flow rate was 0.2 ml/min. A C-18 column (150 × 2.1 mm, 3.5 µm, Agilent, USA) was used for all analysis. The following gradient was used: 5% B at 0 min, 5% B at 5 min, 100% B at 8 min, 100% B at 9 min, 5% B at 18 min and 5% B at 20 min.

The positive ion dataset was run as two batches, with the pooled QC sample repeated once every sixth sample. This produced final spectra as follows: 30 class A; 28

class B; 30 class C; 30 class D; 30 class E; 30 class F; and 39 QC samples spread throughout. The negative ion dataset was run as five batches, with the pooled QC sample injected five times at the start of the run, three times at the end, and once every five samples. The negative ion data has final spectral numbers of: 28 class A; 26 class B; 27 class C; 29 class D; 26 class E; 25 class F; and 66 QC samples spread throughout. Additionally, three blank samples were injected before the starting QC of each run for baseline stabilisation of the LC–MS system – these blank spectra are not included in these published data. Discrepancies with sample numbers are due to some samples being used up entirely as repeats were performed to correct for failed spectra. The positive and negative ion mode LC–MS data were collected in two batches and five batches, respectively, due to an observation that the negative ion responses decreased faster than positive ion responses.

#### Shotgun lipidomics and carnitine profiling spectral acquisition

Lipidomics analysis was carried out as previously described, with slight modifications to previously reported protocols [5,6]. Plasma samples were thawed and extracted, with 10 µL of plasma placed in 1 mL glass vials. Eight µL of an internal standards mixture consisting of ceramide (Cer, d18:1/17:0), phosphoethanolamine (PE, 17:0), phosphatidylserine (PS, 17:0), 1- $\alpha$ -phosphatidylinositol (Soy PI), phosphatidylcholine (PC, 18:3), and phosphoglycerol (PG, 17:0), with a concentration of 10 µM for each lipid, was added. Individual lipids were purchased from Avanti Polar Lipids (Alabaster, AL). Following this, 800 µL of chloroform/methanol/water (20:10:1, V/V/V) containing 300 mM ammonium acetate was added and vortexed. After 10 min, centrifugation at 14000 *g* and 10°C (Eppendorf, Hamburg, Germany) was performed and 100 µL of the lower organic layer was diluted with 100 µL of methanol/chloroform (2:1, V/V) containing 10 mM ammonium acetate.

The final mixture was transferred into a new vial and frozen at  $-20^{\circ}\text{C}$  until required.

Lipidome and carnitine profiling were detected by multiplexed precursor ion scanning (MPIS) analysis. One hundred  $\mu\text{L}$  of final extract was loaded into 96 well plates (Eppendorf, Hamburg, Germany) and sealed with aluminium foil. The final extracts were analysed using a QTRAP<sup>®</sup> 5500 System equipped with a Shimadzu Prominence HPLC system (Shimadzu, Japan). Liquid chromatography and flow injection analysis–mass spectrometry were employed to acquire the data. The electrospray source was operated in the positive ion mode, with optimised source parameters: ionSpray Voltage 5.5 kV, source temperature  $300^{\circ}\text{C}$ , curtain gas 30. Collision energies ranged from 25–30 V for various lipid species in precursor ion or neutral loss scan modes. Flow injection analysis was performed with a mixture of chloroform-methanol-10 mM ammonium acetate in water (300/665/35, V/V/V) at a flow rate of 200  $\mu\text{L}/\text{min}$ . Total analysis time per sample was 3.6 min. PE, PC, PS, PI, PG, Cer and carnitines were monitored by consecutive positive ion mode PIS  $m/z$  141.0, 184.1, 185.0, 277.0, 189.0, 264.2 and 85.1, respectively. Acquired spectra were processed by LipidView<sup>™</sup> and ChemoView<sup>™</sup> software (AB Sciex) for isotope correction, identification and quantification of detected lipid and carnitine species as previously described [5–7].

This lipidomics dataset was run as two batches, with the pooled QC sample injected three times at the start, three times at the end and once every 20 samples. This dataset comprises: 29 class A; 29 class B; 30 class C; 30 class D; 30 class E; 29 class F; 14 QC samples throughout.

### Potential uses

On top of the already published work characterising metabolic phenotype variations during pregnancy, these data also provide a useful baseline and reference dataset to be compared to specific conditions (e.g. hypertensive disorders, diabetes mellitus) in pregnancy. The data were observed to be of good quality via plotting (PCA scores) of QCs alongside samples and noting that the QC samples were tightly clustered. This was observed after processing, full details of which can be found in the previously published research article [1]. Although the positive ion, non-targeted dataset was not randomised during LC–MS collection, the inclusion of QC samples throughout the collection process arguably allows for correction of drift and other systematic noise (e.g. biases correlated with analysis order and/or sample preparation order). The other two datasets were fully pseudo-randomised. The large scale of this dataset and the inclusion of quality control measures makes it very useful for algorithm optimisation and development,

especially those that deal with quality control issues such as batch and drift correction [8–9].

The lipidomics and carnitine dataset ought to be of particular use as a biological reference when studying abnormal pregnancy due to its ability to describe nutrition and energy levels in the subject.

### Availability of supporting data and materials

Supporting data and corresponding ISA-TAB metadata are available in the MetaboLights database [MTBLS146], as well as the *GigaScience* GigaDB repository [10].

### Additional file

**Additional file 1: A detailed, graphical depiction of batch numbers, run order and distribution of QCs for all datasets.**

### Abbreviations

BMI: Body mass index; LC: Liquid chromatography; MS: Mass spectrometry; PC: Phosphatidylcholine; PE: Phosphoethanolamine; PS: Phosphatidylserine; PI: Phosphatidylinositol; PG: Phosphoglycerol; Cer: Ceramide.

### Competing interests

The authors declare that they have no competing interests.

### Authors' contributions

HL, NM, XC and WR performed the experiments; PL collected the clinical samples; HL and JF performed the statistical analysis. HL, HJ, XX, ZC and JW conceived of the study, and participated in its design and coordination. All authors read and approved the final manuscript.

### Acknowledgments

We thank the *GigaScience* team for editing the manuscript. Data were curated during the BBSRC-funded hackathon by the *GigaScience*, ISA-TAB and MetaboLights teams (Grant No. BB/J020265/1). This work was supported by grants from the National Science Foundation of China (Grant No. 1072701) and National Key Basic Research Development Program (Grant No. 2013CB945204).

### Author details

<sup>1</sup>Department of Chemistry, Hong Kong Baptist University, 224 Waterloo Road, Kowloon Tong, Hong Kong, China. <sup>2</sup>BGI-Shenzhen, Building No.11, Beishan Industrial Zone, Yantian District, Shenzhen 518083, China. <sup>3</sup>Department of Biology, University of Copenhagen, Ole Maaløes Vej 5, DK-2200 Copenhagen, Denmark.

Received: 29 November 2014 Accepted: 11 March 2015

Published online: 09 April 2015

### References

- Luan H, Meng N, Liu P, Feng Q, Lin S, Fu J, et al. Pregnancy-induced metabolic phenotype variations in maternal plasma. *J Proteome Res*. 2014;13(3):1527–36.
- Luan H, Chen X, Zhong S, Yuan X, Meng N, Zhang J, et al. Serum metabolomics reveals lipid metabolism variation between coronary artery disease and congestive heart failure: a pilot study. *Biomarkers*. 2013;18(4):314–21.
- Zelena E, Dunn WB, Broadhurst D, Francis-McIntyre S, Carroll KM, Begley P, et al. Development of a robust and repeatable UPLC–MS method for the long-term metabolomic study of human serum. *Anal Chem*. 2009;81(4):1357–64.
- Want EJ, Wilson ID, Gika H, Theodoridis G, Plumb RS, Shockcor J, et al. Global metabolic profiling procedures for urine using UPLC–MS. *Nat Protoc*. 2010;5(6):1005–18.
- Ejsing CS, Sampaio JL, Surendranath V, Duchoslav E, Ekroos K, Klemm RW, et al. Global analysis of the yeast lipidome by quantitative shotgun mass spectrometry. *Proc Natl Acad Sci U S A*. 2009;106(7):2136–41.

6. Yang Q, Manicke NE, Wang H, Petucci C, Cooks RG, Ouyang Z. Direct and quantitative analysis of underivatized acylcarnitines in serum and whole blood using paper spray mass spectrometry. *Anal Bioanal Chem.* 2012;404(5):1389–97.
7. Wang-Sattler R, Yu Z, Herder C, Messias AC, Floegel A, He Y, et al. Novel biomarkers for pre-diabetes identified by metabolomics. *Mol Syst Biol.* 2012;8:615.
8. Sumner LW, Amberg A, Barrett D, Beale MH, Beger R, Daykin CA, et al. Proposed minimum reporting standards for chemical analysis Chemical Analysis Working Group (CAWG) Metabolomics Standards Initiative (MSI). *Metabolomics.* 2007;3(3):211–21.
9. Salek RM, Steinbeck C, Viant MR, Goodacre R, Dunn WB. The role of reporting standards for metabolite annotation and identification in metabolomic studies. *Gigascience.* 2013;2(1):13.
10. Luan H, Meng N, Liu P, Feng Q, Lin S, Fu J, et al. Nontargeted metabolomics and lipidomics HPLC-MS data from maternal plasma of 180 healthy pregnant women. *GigaScience Database.* 2015. <http://dx.doi.org/10.5524/100108>.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

